

A PhD thesis I would like to do again.

Lucas Pastor



Definition


A graph G is **perfect** if every induced subgraph H of G satisfies $\chi(H) = \omega(H)$.

Definition

A graph G is **perfect** if every induced subgraph H of G satisfies $\chi(H) = \omega(H)$.

Definition

A graph G is **claw-free** if it does not contain any induced subgraph isomorphic to $K_{1,3}$ (the claw).




claw

Definition

Given a graph G and a set $L(v)$ of colors for each vertex v , we say that G is ***L-colorable*** if we can find a coloring c such that $c(v) \in L(v)$ for all $v \in V(G)$.


Definition

Given a graph G and a set $L(v)$ of colors for each vertex v , we say that G is *L-colorable* if we can find a coloring c such that $c(v) \in L(v)$ for all $v \in V(G)$.



Definition

Given a graph G and a set $L(v)$ of colors for each vertex v , we say that G is *L-colorable* if we can find a coloring c such that $c(v) \in L(v)$ for all $v \in V(G)$.



Definition

Given an integer k , a graph G is k -**choosable** if it is L -colorable for every assignment L that satisfies $|L(v)| = k$ for all $v \in V(G)$.

Definition

Given an integer k , a graph G is k -**choosable** if it is L -colorable for every assignment L that satisfies $|L(v)| = k$ for all $v \in V(G)$.

Definition

The **choice number**, or **list-chromatic number** $\chi_\ell(G)$ of G is the smallest k such that G is k -choosable.

Definition

Given an integer k , a graph G is k -**choosable** if it is L -colorable for every assignment L that satisfies $|L(v)| = k$ for all $v \in V(G)$.

Definition

The **choice number**, or **list-chromatic number** $\chi_\ell(G)$ of G is the smallest k such that G is k -choosable.

Conjecture Vizing 1975

For every graph G , $\chi_\ell(\mathcal{L}(G)) = \chi(\mathcal{L}(G))$.

Definition

Given an integer k , a graph G is k -**choosable** if it is L -colorable for every assignment L that satisfies $|L(v)| = k$ for all $v \in V(G)$.

Definition

The **choice number**, or **list-chromatic number** $\chi_\ell(G)$ of G is the smallest k such that G is k -choosable.

Conjecture Vizing 1975

For every graph G , $\chi_\ell(\mathcal{L}(G)) = \chi(\mathcal{L}(G))$.

Conjecture Gravier, Maffray 1997

For every claw-free graph G , $\chi_\ell(G) = \chi(G)$.

Conjecture

For every claw-free graph G , $\chi_\ell(G) = \chi(G)$.

Conjecture

For every claw-free graph G , $\chi_\ell(G) = \chi(G)$.

We are interested in the case where G is *perfect*.

Conjecture

For every claw-free graph G , $\chi_\ell(G) = \chi(G)$.

We are interested in the case where G is *perfect*.

Theorem

Let G be a claw-free perfect graph $\omega(G) \leq 4$. Then $\chi_\ell(G) = \chi(G)$.

Conjecture

For every claw-free graph G , $\chi_\ell(G) = \chi(G)$.

We are interested in the case where G is **perfect**.

Theorem

Let G be a claw-free perfect graph $\omega(G) \leq 4$. Then $\chi_\ell(G) = \chi(G)$.


We achieved that thanks to a decomposition theorem of Chátal and Sbihi **and** a structural theorem of Maffray and Reed.

Theorem Chvátal, Sbihi 1988

Every claw-free perfect graph either has a *clique-cutset*, or is a *peculiar graph*, or is an *elementary graph*.


Theorem Chvátal, Sbihi 1988

Every claw-free perfect graph either has a *clique-cutset*, or is a *peculiar graph*, or is an *elementary graph*.




Theorem Chvátal, Sbihi 1988

Every claw-free perfect graph either has a *clique-cutset*, or is a *peculiar graph*, or is an *elementary graph*.




Theorem Chvátal, Sbihi 1988

Every claw-free perfect graph either has a **clique-cutset**, or is a **peculiar graph**, or is an **elementary graph**.



Peculiar graphs



clique



at least one non-edge



complete adjacency

Lemma

Let G be a peculiar graph with $\omega(G) \leq 4$ (*unique* in this case). Then, $\chi_\ell(G) = \chi(G)$.

Lemma

Let G be a peculiar graph with $\omega(G) \leq 4$ (*unique* in this case). Then, $\chi_\ell(G) = \chi(G)$.

Proof


- If some pair of non-adjacent vertices u, v share a color, let $\alpha = c(u) = c(v)$ and we can easily color $G - \{u, v\}$ without using α .
- If no such pair exists, we can find a coloring by Hall's theorem (we have enough color to directly color G).

Theorem

A graph G is elementary if and only if it is an augmentation of a the line-graph H of a bipartite multigraph B .


Theorem

A graph G is elementary if and only if it is an augmentation of a the line-graph H of a bipartite multigraph B .




Theorem

A graph G is elementary if and only if it is an augmentation of a the line-graph H of a bipartite multigraph B .




Theorem

A graph G is elementary if and only if it is an augmentation of a the line-graph H of a bipartite multigraph B .




Theorem

A graph G is elementary if and only if it is an augmentation of a the line-graph H of a bipartite multigraph B .



Theorem

A graph G is elementary if and only if it is an augmentation of a the line-graph H of a bipartite multigraph B .



Lemma

Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_\ell(G) = \chi(G)$.

Lemma

Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_\ell(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.

List-coloring claw-free perfect graphs


Lemma

Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_\ell(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.



List-coloring claw-free perfect graphs


Lemma

Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_{\ell}(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.



List-coloring claw-free perfect graphs


Lemma

Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_\ell(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.





Lemma

Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_\ell(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.




Lemma

Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_{\ell}(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.



Lemma


Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_{\ell}(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.

L -colorable \iff the color changes



Lemma


Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_{\ell}(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.

L -colorable \iff the color changes



List-coloring claw-free perfect graphs


Lemma

Let G be an elementary graph with $\omega(G) \leq 4$. Then $\chi_\ell(G) = \chi(G)$.

Proof

By induction on h , the number of augmented flat edges :

- If $h = 0$, use Gavlin's theorem (LCC is true for line-graphs of bipartite multigraphs).
- If $h > 0$, we use a gadget.



Proof of the main theorem

The graph G is decomposed by clique cutset into:

- peculiar graphs
- elementary graphs.

Proof of the main theorem

The graph G is decomposed by clique cutset into:

- peculiar graphs ✓
- elementary graphs.

Proof of the main theorem

The graph G is decomposed by clique cutset into:

- peculiar graphs ✓
- elementary graphs. ✓

Proof of the main theorem

The graph G is decomposed by clique cutset into:

- peculiar graphs ✓
- elementary graphs. ✓

We still need to glue back all pieces!

Proof of the main theorem

The graph G is decomposed by clique cutset into:

- peculiar graphs ✓
- elementary graphs. ✓

We still need to glue back all pieces!

Within the list-coloring context, clique cutsets are not as convenient as for the classical coloring.

Proof of the main theorem

The graph G is decomposed by clique cutset into:

- peculiar graphs ✓
- elementary graphs. ✓

We still need to glue back all pieces!

Within the list-coloring context, clique cutsets are not as convenient as for the classical coloring.

We still manage to deal with them by using **Galvin's theorem**.

Theorem *Garey, Johnson, Stockmeyer 1974*

Deciding whether a graph is k -colorable is NP-complete for each $k \geq 3$.

Theorem Garey, Johnson, Stockmeyer 1974

Deciding whether a graph is k -colorable is NP-complete for each $k \geq 3$.

As it is well known, the k -coloring problem is polynomial for perfect graphs.

But let us take a look at **other graph classes**.

Theorem Kamiński, Lozin 2007

For any fixed $k, g \geq 3$, the k -coloring problem is NP-complete in the class of graphs with **girth** at least g .

girth : length of the shortest cycle.

Theorem Kamiński, Lozin 2007

For any fixed $k, g \geq 3$, the k -coloring problem is NP-complete in the class of graphs with **girth** at least g .

girth : length of the shortest cycle.

Theorem Holyer 1981

For any fixed $k \geq 3$, the k -coloring problem is NP-complete in the class of H -free graphs where H contains a claw.

Theorem Kamiński, Lozin 2007

For any fixed $k, g \geq 3$, the k -coloring problem is NP-complete in the class of graphs with **girth** at least g .

girth : length of the shortest cycle.

Theorem Holyer 1981

For any fixed $k \geq 3$, the k -coloring problem is NP-complete in the class of H -free graphs where H contains a claw.

By combining these two theorems, we have the following :

Corollary

For any fixed $k \geq 3$ and H a forbidden induced subgraph that is **not a collection of paths**, deciding whether a H -free graph is k -colorable is NP-complete.

What was known?

k -coloring of P_ℓ -free graphs.

What was known?

k -coloring of P_ℓ -free graphs.

$\ell \setminus k$	≤ 2	3	4	≥ 5
≤ 4	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>
5	<i>P</i>	<i>P</i>	<i>P</i>	<i>P</i>
6	<i>P</i>	<i>P</i>	?	<i>NPC</i>
7	<i>P</i>	<i>P</i>	<i>NPC</i>	<i>NPC</i>
≥ 8	<i>P</i>	?	<i>NPC</i>	<i>NPC</i>

...

⋮

What was known?

k -coloring of P_ℓ -free graphs.

$\ell \setminus k$	≤ 2	3	④	≥ 5
≤ 4	P	P	P	P
5	P	P	P	P
⑥	P	P	⑦	NPC
7	P	P	NPC	NPC
≥ 8	P	?	NPC	NPC

...

⋮

What is known?

k -coloring of P_ℓ -free graphs.

$\ell \setminus k$	≤ 2	3	4	≥ 5
≤ 4	P	P	P	P
5	P	P	P	P
6	P	P	P	NPC
7	P	P	NPC	NPC
≥ 8	P	?	NPC	NPC

...


⋮

Theorem Chudnovsky, Spirkl, Zhong 2018

There exists a polynomial time algorithm for the 4-coloring problem for P_6 -free graphs.

Theorem Maffray, Pastor

There is polynomial time algorithm that determines whether a (P_6, bull) -free graph is 4-colorable, and if it is, produces a 4-coloring.




Definition

A **homogeneous set** is a set $S \subseteq V(G)$ such that every vertex in $V(G) \setminus S$ is either complete to S or anti-complete to S .


Definition

A **homogeneous set** is a set $S \subseteq V(G)$ such that every vertex in $V(G) \setminus S$ is either complete to S or anti-complete to S .



Definition

A **homogeneous set** is a set $S \subseteq V(G)$ such that every vertex in $V(G) \setminus S$ is either complete to S or anti-complete to S .



Definition

Quasi-prime graph A graph G is **quasi-prime** if every non-trivial homogeneous set of G is a clique.

Lemma


It is sufficient to produce a 4-coloring for any (P_6, bull) -free graph G that satisfies the following properties :

1. G is K_5 -free and double-wheel-free.
2. G and \overline{G} are connected.
3. G is quasi-prime.

Lemma

It is sufficient to produce a 4-coloring for any (P_6, bull) -free graph G that satisfies the following properties :


1. G is K_5 -free and double-wheel-free.
2. G and \overline{G} are connected.
3. G is quasi-prime.



double-wheel

Proof of G and \overline{G} connected.


Proof of G and \overline{G} connected.



G


If G is not connected.

Proof of G and \overline{G} connected.




If G is not connected.

Proof of G and \overline{G} connected.




If G is not connected.

Proof of G and \overline{G} connected.



If G is not connected.
Keep the maximum over all $\chi(G_i)$.


Proof of G and \overline{G} connected.



G


If \overline{G} is not connected.

Proof of G and \overline{G} connected.



If \overline{G} is not connected.


Proof of G and \overline{G} connected.



If \overline{G} is not connected.

Test 3-colorability of each co-components with known algorithms.

Proof of G and \overline{G} connected.



If \overline{G} is not connected.

Test 3-colorability of each co-components with known algorithms.

Refine to test whether they are 1-, 2- or 3-colorable.

Lemma


Let G be a quasi-prime bull-free graph that contains no K_5 and no double-wheel. Then at least one of the following holds :

1. G is gem-free.
2. G contains a *magnet*.
3. G contains the special graph.


Lemma

Let G be a quasi-prime bull-free graph that contains no K_5 and no double-wheel. Then at least one of the following holds :

1. G is gem-free.
2. G contains a *magnet*.
3. G contains the special graph.



gem



special graph

Lemma

Let G be a $(P_6, \text{bull}, \text{gem})$ -free graph, then G is either :

1. Perfect.
2. Of bounded clique-width.

Lemma

Let G be a $(P_6, \text{bull}, \text{gem})$ -free graph, then G is either :

1. Perfect.
2. Of bounded clique-width.


1. Algorithm for bull-free perfect graphs.
2. Courcelle's theorem.

Definition

A subgraph F of G is a **magnet** if every vertex of $G \setminus F$ has two neighbours $u, v \in V(F)$ such that $uv \in E(F)$.


Definition

A subgraph F of G is a **magnet** if every vertex of $G \setminus F$ has two neighbours $u, v \in V(F)$ such that $uv \in E(F)$.




Definition

A subgraph F of G is a **magnet** if every vertex of $G \setminus F$ has two neighbours $u, v \in V(F)$ such that $uv \in E(F)$.




Definition

A subgraph F of G is a **magnet** if every vertex of $G \setminus F$ has two neighbours $u, v \in V(F)$ such that $uv \in E(F)$.



Definition

A subgraph F of G is a **magnet** if every vertex of $G \setminus F$ has two neighbours $u, v \in V(F)$ such that $uv \in E(F)$.



Lemma

Let G be a quasi-prime bull-free graph that contains no K_5 and no double-wheel. Then at least one of the following holds:

- G is gem-free.
- G contains a magnet.
- G contains the special graph.

Lemma

Let G be a quasi-prime bull-free graph that contains no K_5 and no double-wheel. Then at least one of the following holds:

- G is gem-free. ✓
- G contains a magnet.
- G contains the special graph.

Lemma

Let G be a quasi-prime bull-free graph that contains no K_5 and no double-wheel. Then at least one of the following holds:


- G is gem-free. ✓
- G contains a magnet. ✓
- G contains the special graph.

Lemma


Let G be a quasi-prime bull-free graph that contains no K_5 and no double-wheel. Then at least one of the following holds:

- G is gem-free. ✓
- G contains a magnet. ✓
- G contains the special graph.


Coloring (P_6, bull) -free graphs



Coloring (P_6, bull) -free graphs




Coloring (P_6 , bull)-free graphs



Coloring (P_6 , bull)-free graphs

2-list-coloring



Theorem Maffray, Pastor

There is a polynomial time algorithm that determines whether a (P_6, bull) -free graphs is 4-colorable, and if it is, produces a 4-coloring.

Theorem Maffray, Pastor

There is a polynomial time algorithm that determines whether a (P_6, bull) -free graph is 4-colorable, and if it is, produces a 4-coloring.

Theorem Maffray, Pastor

For any fixed k , there is a polynomial algorithm that determines if a $(P_6, \text{bull}, \text{gem})$ -free graph is k -colorable and if it is, produces a k -coloring.

Conclusion

- At the beginning of my PhD, I realised I knew close to nothing, and he knew **A LOT**.
- During my PhD thesis, I participated in 6 (3 with only the two of us) papers where Frédéric was also a co-author.
- These 3 years of PhD were a walk in the park thanks to him.

Conclusion

- At the beginning of my PhD, I realised I knew close to nothing, and he knew **A LOT**.
- During my PhD thesis, I participated in 6 (3 with only the two of us) papers where Frédéric was also a co-author.
- These 3 years of PhD were a walk in the park thanks to him.

Thank you for your attention.